

1N-46

115135

P.26

NASA  
Reference  
Publication  
1275

August 1992

# SAGE I Data User's Guide

Leonard R. McMaster,  
William P. Chu,  
and Michael W. Rowland

(NASA-RP-1275) SAGE I DATA USER'S  
GUIDE (NASA) 26 p

N92-33097

Unclass

H1/46 0115135

NASA

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200
201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300
301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400
401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500
501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600
601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700
701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800
801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900
901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000
1001	1002	1003	1004	1005	1006	1007	1008	1009	1010	1011	1012	1013	1014	1015	1016	1017	1018	1019	1020	1021	1022	1023	1024	1025	1026	1027	1028	1029	1030	1031	1032	1033	1034	1035	1036	1037	1038	1039	1040	1041	1042	1043	1044	1045	1046	1047	1048	1049	1050	1051	1052	1053	1054	1055	1056	1057	1058	1059	1060	1061	1062	1063	1064	1065	1066	1067	1068	1069	1070	1071	1072	1073	1074	1075	1076	1077	1078	1079	1080	1081	1082	1083	1084	1085	1086	1087	1088	1089	1090	1091	1092	1093	1094	1095	1096	1097	1098	1099	1100
1101	1102	1103	1104	1105	1106	1107	1108	1109	1110	1111	1112	1113	1114	1115	1116	1117	1118	1119	1120	1121	1122	1123	1124	1125	1126	1127	1128	1129	1130	1131	1132	1133	1134	1135	1136	1137	1138	1139	1140	1141	1142	1143	1144	1145	1146	1147	1148	1149	1150	1151	1152	1153	1154	1155	1156	1157	1158	1159	1160	1161	1162	1163	1164	1165	1166	1167	1168	1169	1170	1171	1172	1173	1174	1175	1176	1177	1178	1179	1180	1181	1182	1183	1184	1185	1186	1187	1188	1189	1190	1191	1192	1193	1194	1195	1196	1197	1198	1199	1200
1201	1202	1203	1204	1205	1206	1207	1208	1209	1210	1211	1212	1213	1214	1215	1216	1217	1218	1219	1220	1221	1222	1223	1224	1225	1226	1227	1228	1229	1230	1231	1232	1233	1234	1235	1236	1237	1238	1239	1240	1241	1242	1243	1244	1245	1246	1247	1248	1249	1250	1251	1252	1253	1254	1255	1256	1257	1258	1259	1260	1261	1262	1263	1264	1265	1266	1267	1268	1269	1270	1271	1272	1273	1274	1275	1276	1277	1278	1279	1280	1281	1282	1283	1284	1285	1286	1287	1288	1289	1290	1291	1292	1293	1294	1295	1296	1297	1298	1299	1300
1301	1302	1303	1304	1305	1306	1307	1308	1309	1310	1311	1312	1313	1314	1315	1316	1317	1318	1319	1320	1321	1322	1323	1324	1325	1326	1327	1328	1329	1330	1331	1332	1333	1334	1335	1336	1337	1338	1339	1340	1341	1342	1343	1344	1345	1346	1347	1348	1349	1350	1351	1352	1353	1354	1355	1356	1357	1358	1359	1360	1361	1362	1363	1364	1365	1366	1367	1368	1369	1370	1371	1372	1373	1374	1375	1376	1377	1378	1379	1380	1381	1382	1383	1384	1385	1386	1387	1388	1389	1390	1391	1392	1393	1394	1395	1396	1397	1398	1399	1400
1401	1402	1403	1404	1405	1406	1407	1408	1409	1410	1411	1412	1413	1414	1415	1416	1417	1418	1419	1420	1421	1422	1423	1424	1425	1426	1427	1428	1429	1430	1431	1432	1433	1434	1435	1436	1437	1438	1439	1440	1441	1442	1443	1444	1445	1446	1447	1448	1449	1450	1451	1452	1453	1454	1455	1456	1457	1458	1459	1460	1461	1462	1463	1464	1465	1466	1467	1468	1469	1470	1471	1472	1473	1474	1475	1476	1477	1478	1479	1480	1481	1482	1483	1484	1485	1486	1487	1488	1489	1490	1491	1492	1							

**NASA  
Reference  
Publication  
1275**

1992

# SAGE I Data User's Guide

Leonard R. McMaster  
and William P. Chu  
*Langley Research Center  
Hampton, Virginia*

Michael W. Rowland  
*ST Systems Corporation (STX)  
Hampton, Virginia*



National Aeronautics and  
Space Administration  
Office of Management  
Scientific and Technical  
Information Program



## Preface

The data products from the Stratospheric Aerosol and Gas Experiment I (SAGE I), flown on the Applications Explorer Mission 2 (AEM-2) satellite, have been archived on magnetic tapes at the National Space Sciences Data Center, NASA Goddard Space Flight Center, Greenbelt, Maryland 20771, and are now available to researchers. The AEM-2 satellite was launched on February 18, 1979, but because of the failure of the spacecraft power subsystem, the mission was terminated on November 11, 1981. Over the operational life of the spacecraft, the instrument collected 33 months of scientific data, including vertical profiles of stratospheric aerosol, ozone, and nitrogen dioxide. This document is intended to serve as a guide for using these data sets in scientific investigations of stratospheric chemistry related to aerosol, ozone, nitrogen dioxide, dynamics, and climate change. There are brief descriptions of the instrument operation; data collection, processing, and validation; and some descriptions of the scientific analyses that have been conducted. The SAGE I data products, containing the aerosol extinction data at 1000 and 450 nm, the ozone concentration data, and the nitrogen dioxide concentration data, are described in detail in appendixes A, B, and C.

Over the years, the SAGE I experiment's development and research have been guided by the SAGE I Science Team, made up of the following people:

M. P. McCormick, Principal Investigator, NASA Langley Research Center (LaRC); R. Craig, Florida State University; D. M. Cunnold and G. W. Grams, Georgia Institute of Technology; B. M. Herman, University of Arizona; M. Hirono, Kyoto University; D. E. Miller, British Meteorological Office; D. G. Murcray, University of Denver; T. J. Pepin, University of Wyoming; W. G. Planet, National Oceanic and Atmospheric Administration National Meteorological Center; and P. B. Russell, NASA Ames Research Center.

The efforts of the LaRC SAGE I Data Processing Team are also gratefully acknowledged.



## Contents

Preface . . . . .	iii
Summary . . . . .	1
Introduction . . . . .	1
AEM-2 SAGE I Mission . . . . .	1
Instrument Description and Operation . . . . .	2
Data Processing . . . . .	3
Data Validation . . . . .	6
Data Products . . . . .	6
Scientific Studies . . . . .	8
Concluding Remarks . . . . .	9
Appendix A—SAGE I Aerosol Profile Record Format . . . . .	10
Appendix B—SAGE I Ozone Profile Record Format . . . . .	13
Appendix C—SAGE I Nitrogen Dioxide Profile Record Format . . . . .	16
Appendix D—Meteorological Data Notes . . . . .	19
References . . . . .	20





## Summary

The Stratospheric Aerosol and Gas Experiment I (SAGE I) instrument was launched on board the Applications Explorer Mission-2 (AEM-2) satellite, which was dedicated to the SAGE I mission. SAGE I provided global data collection of aerosol vertical extinction profiles, ozone vertical concentration profiles, and nitrogen dioxide vertical concentration profiles. The SAGE I instrument was a Sun photometer that measured the attenuation of solar radiation in four spectral regions through the Earth's atmosphere during spacecraft sunrise and sunset. The solar radiance data were combined with spacecraft ephemeris and National Oceanic and Atmospheric Administration (NOAA) meteorological data and then numerically inverted to yield altitude profiles of aerosol extinction at wavelengths of 1000 and 450 nm and altitude profiles of ozone and nitrogen dioxide concentration. The SAGE I aerosol data were validated by comparison with correlative lidar and dustsonde in situ measurements, the ozone data were validated by comparison with balloon electrochemical cell (ECC) ozonesonde and rocket measurements, and the nitrogen dioxide measurements were compared with climatology. These data are currently archived at the National Space Sciences Data Center (NSSDC). This publication describes the SAGE I experiment, instrument characteristics, and mode of operation; outlines the method of the data inversion; and explains to potential users the format of the data products archived at NSSDC. Results of the data validation and examples of data products are also presented to demonstrate the quality of the data and some of its applications in atmospheric studies.

## Introduction

SAGE I was the second in a series of satellite experiments using the solar occultation technique to monitor and study stratospheric trace constituents. The Stratospheric Aerosol Measurement II (SAM II), the first automatic instrument to use this technique from space, was launched October 24, 1978, on the Nimbus 7 spacecraft (McCormick et al. 1979). The SAM II instrument uses a single spectral channel centered at a wavelength of 1000 nm for monitoring stratospheric aerosols. Because of the Nimbus 7's Sun-synchronous orbit, SAM II collects stratospheric aerosol data exclusively in the two polar regions at latitudes extending from 64° to 80° N and from 64° to 80° S. At the time this paper was prepared, SAM II was still operational and in its 13th year of data collection.

The Stratospheric Aerosol and Gas Experiment I (SAGE I) was launched February 18, 1979, aboard

the Applications Explorer Mission-2 (AEM-2) satellite (McCormick et al. 1979). The SAGE I instrument had four spectral channels centered at wavelengths of 1000, 600, 450, and 385 nm for measuring the atmospheric extinction on account of aerosols, ozone, and nitrogen dioxide. The AEM-2 satellite was placed in an orbit of approximately 600 km at an inclination of 56° to extend the latitudinal coverage for the solar occultation measurements from 79° S to 79° N. The SAGE I instrument collected data for almost 3 years until the AEM-2 satellite power subsystem failed. The processed SAGE I data have been archived at the National Space Sciences Data Center (NSSDC), NASA Goddard Space Flight Center, Greenbelt, Maryland 20771.

The most recent of these solar occultation experiments, the Stratospheric Aerosol and Gas Experiment II (SAGE II), uses seven spectral channels for measuring the atmospheric extinction of aerosols, ozone, nitrogen dioxide, and water vapor and is in an orbit similar to that of SAGE I, which allows data collection at latitudes from 80° S to 80° N. The SAGE II instrument was launched October 5, 1984, aboard the Earth Radiation Budget Satellite (ERBS) and was still collecting data after more than 7 years of operation at the time this paper was prepared (McMaster 1986). As the SAGE II data are processed and validated, they are also being archived at the NSSDC.

## AEM-2 SAGE I Mission

The scientific objective of SAGE I was to develop a global stratospheric aerosol, ozone, and nitrogen dioxide data base that could be used for the investigation of the spatial and temporal variations of these species caused by seasonal and short-term meteorological variations, atmospheric chemistry and microphysics, and transient phenomena such as volcanic eruptions. The data base could also be used for the study of trends, atmospheric dynamics and transport, and potential climatic effects. The SAGE I sensor was designed to measure the attenuation of solar radiation resulting from atmospheric aerosol, ozone, and nitrogen dioxide at four spectral regions through the Earth's atmosphere during each spacecraft sunrise and sunset (fig. 1). The SAGE I sensor was a dedicated instrument on the AEM-2 satellite which, had its orbit tailored to maximize geographic coverage for solar occultation sampling. The AEM-2 satellite, with its orbital altitude of approximately 600 km and angle of inclination of 56°, provided 15 sunrise and 15 sunset measurements each day, with successive sunrise or sunset measurements occurring about 25° of longitude apart. The latitude of the sunrise or

sunset measurements varied slowly from day to day and provided near-global coverage every 3 to 4 weeks. As the seasons changed, the maximum latitude of the SAGE I measurements ranged from  $79^{\circ}$  S to  $79^{\circ}$  N. Figure 2 shows the latitudinal coverage of the SAGE I mission over the 33 months of instrument operation. Because of the spacecraft power subsystem problems that reduced the power capacity of its batteries, sunrise measurements from SAGE I were terminated after August 1979.






Figure 1. Solar occultation measurement geometry.

## Instrument Description and Operation


The SAGE I instrument was a four-channel Sun photometer that used a Cassegrainian telescope, holographic grating, and four silicon photodiodes to define the four-spectral-channel bandpass. Figure 3 is a schematic of the SAGE I instrument. Solar radiation was reflected off a scan mirror into the telescope, with an image of the Sun being formed at the focal plane. The instrument's instantaneous field of view, defined by the aperture on the focal plane, was a 30-arcsec circle that produced a vertical resolution at the tangent point of about 0.5 km. Radiation passing through the aperture was transferred to the spectrometer section of the instrument which contained the holographic grating and four separate detector systems. The holographic grating dispersed the incoming radiation into the four spectral regions centered at wavelengths of 1000, 600, 450, and 385 nm. Slits on the Rowland circle of the grating defined the spectral bandpass of the four spectral channels. The bandpasses were 50, 30, 20, and 30 nm, respectively, for the above-mentioned wavelengths. The entire imaging and spectrometer system was inside the azimuth gimbal to allow the instrument to be pointed at the Sun without image rotation. The azimuth gimbal could be rotated over  $360^{\circ}$  so that measurements could be made at any azimuthal angle.



(a) 1979.



(b) 1980.



(c) 1981.

Figure 2. Latitudinal coverage for SAGE I measurements. Pluses indicate sunrise measurements and diamonds indicate sunset measurements.




Figure 3. Schematic of SAGE I instrument. (From McCormick et al. 1979.)

The operation of the instrument, during each sunrise and sunset measurement, was totally automatic. Prior to each sunrise or sunset, the instrument was rotated in azimuth to its predicted solar acquisition position. When the Sun entered the instrument's field of view, the instrument adjusted its azimuth position to lock onto the radiometric center of the Sun to within  $\pm 45$  arcsec and then acquired the Sun by rotating its scan mirror to the proper elevation angle. As the Sun traversed between the horizon and the tangent height of 150 km, the elevation mirror scanned vertically across the solar disk. (See fig. 4.) The two solid lines in the figure represent the top and bottom edges of the solar disk during a sunset event. The gradual shrinking of the vertical shape of the solar image at the lower altitudes was caused by increasing atmospheric refraction. The dashed line represents the relative motion of the instrument elevation mirror as it scanned the Sun at a nominal rate of 15 arcmin/sec. The radiometric channel data were sampled at a rate of 64 samples per second per channel, digitized to 12-bit resolution, and recorded for later transmission back to Earth. Additional SAGE I instrument information can be found in McCormick et al. (1979).




Figure 4. Typical SAGE I elevation scanning of solar disk for a sunset measurement event.

## Data Processing

The SAGE I measurement data were processed with the algorithm developed by Chu and McCormick (1979) and its updated revision as discussed by Chu et al. (1989). In this section, a brief outline of the SAGE I algorithm is presented.

The measured radiance from the SAGE I instrument can be related to the atmospheric optical properties with the Lambert-Beer relation:

$$H_{\lambda}(t) = \int F(\theta, \phi) S(\theta, \phi, t) \exp[-\tau_{\lambda}(\theta)] d\Omega \quad (1)$$

where

$H_{\lambda}(t)$	measured radiance at time $t$ for channel wavelength $\lambda$
$F(\theta, \phi)$	instrumental field-of-view function
$S(\theta, \phi, t)$	extraterrestrial solar radiance profile within the SAGE I spectral bandwidth at time $t$ corrected for atmospheric refraction effects
$\tau_{\lambda}(\theta)$	optical thickness of atmosphere for elevation angle $\theta$ and wavelength $\lambda$
$\phi$	azimuthal angle
$\Omega$	total solid angle

Each elevation angle  $\theta$  corresponds uniquely to an atmospheric tangent height  $h_t$ , the optical thickness  $\tau_{\lambda}(\theta)$  can be related to atmospheric extinction properties through the following equation:

$$\tau_{\lambda}(h_t) = \int [\beta_a(h) + \beta_{O_3}(h) + \beta_{NO_2}(h) + \beta_{Ray}(h)] dh \quad (2)$$

where

$\beta_a(h)$	profile of aerosol extinction versus altitude
--------------	---

$\beta_{\text{O}_3}(h)$	profile of ozone extinction versus altitude
$\beta_{\text{NO}_2}(h)$	profile of nitrogen dioxide extinction versus altitude
$\beta_{\text{Ray}}(h)$	profile of Rayleigh extinction versus altitude
$d\rho$	differential path length through atmosphere

The integral is evaluated from the spacecraft position to the Sun.

The processing of the SAGE I data to generate the various species profiles was done in three steps. First, the measured radiance data were reduced with the spacecraft ephemeris, atmospheric refraction calculation, and the Sun-scan data and put into profiles of limb optical thickness  $\tau_\lambda(h_t)$  as a function of tangent height  $h_t$  in the atmosphere for each channel centered at wavelength  $\lambda$ . The high-altitude solar-scan profiles for each channel were used as calibrated solar-limb profiles out of the atmosphere.

The second step was to subtract the estimated Rayleigh contribution along each limb path for each channel and to separate the four spectral optical thickness profiles into aerosol optical thickness profiles at 1000 and 450 nm, an ozone optical thickness profile at 600 nm, and a nitrogen dioxide optical thickness profile at 385 nm with the method described by Chu and McCormick (1979) and Chu et al. (1989). The Rayleigh contributions were computed from coincident temperature versus height profiles that were provided by the National Oceanic and Atmospheric Administration (NOAA).

The third step was to invert the species' optical thickness profiles into extinction profiles. Through division of the atmosphere into  $N$  homogeneous layers, the integral equation can be reduced to a system of linear equations as follows:

$$\tau_{\alpha i} = \sum_j^N \rho_{ij} \beta_{\alpha j} \quad (3)$$

where


$\tau_{\alpha i}$	measured limb optical thickness at $i$ th layer for species $\alpha$
$\rho_{ij}$	path length of the Sun's ray in $j$ th layer with its tangent height at $i$ th layer
$\beta_{\alpha j}$	averaged extinction coefficient for species $\alpha$ in $j$ th layer

Equation (3) is then inverted with Twomey's modification of Chahine's nonlinear relaxation algorithm as described by Chu and McCormick (1979) and Chu et al. (1989). The SAGE I inversion algorithm generated vertical aerosol extinction profiles at 1000 and 450 nm and ozone and nitrogen dioxide vertical concentration profiles.


An example of the SAGE I aerosol data inversion is shown in figure 5. The aerosol extinction profiles of 1000 nm that are shown are from SAGE I radiance data collected April 24, 1979, at 0545 and 0722 LCT. They illustrate the background aerosol condition versus the presence of volcanic plume from the eruption of Soufrière volcano (McCormick et al. 1982). The error bars shown in the figures are estimates of the random errors associated with the calculations of the aerosol extinction that result from both the radiometric measurement and the mathematical inversion. These errors (also archived at NSSDC along with the constituent profiles) include contributions from random measurement and inversion noise, NOAA temperature uncertainties, and altitude uncertainties. Below an altitude of 25 km, where aerosol extinction exceeds molecular extinction by 50 percent, the total error in the retrieved aerosol extinction coefficient is typically less than 10 percent. Therefore, even under most background or nonvolcanic conditions, the extinction resulting from stratospheric aerosols can be measured to accuracies within 10 percent.

An example of the SAGE I ozone data inversion is shown in figure 6. The dashed-line profile of ozone partial pressure is from SAGE I radiance data collected on April 2, 1979. It illustrates a typical mid-latitude springtime ozone profile. The profile shown by the solid line in the figure is a coincident electrochemical cell (ECC) ozonesonde profile obtained at Garmisch-Partenkirchen, Germany (Reiter and McCormick 1982). The error bars associated with the SAGE I ozone profile are estimates of the random errors from the calculation of the ozone concentration profile. These errors are estimated similar to the errors in the aerosol extinction profiles. Generally, the uncertainties of the ozone concentration profiles retrieved from the SAGE I measurements are less than 10 percent from cloud-top height to a geometric altitude of about 45 km.

An example of the SAGE I nitrogen dioxide data inversion with the 385-nm channel is shown in figure 7. (See also WMO 1981.) Chu and McCormick (1986) suggested that measurement noise and difficulties associated with removing the Rayleigh, ozone, and aerosol contributions from this single wavelength retrieval led to an accuracy of approximately 20 percent, as shown by the error bars. The effect of the



(a) Background conditions. 0722 LCT; long. 85.4° W; lat. 19.8° N.



(b) Perturbed conditions resulting from La Soufrière volcano eruption. 0545 LCT; long. 61.0° W; lat. 19.5° N.

Figure 5. SAGE I aerosol extinction profile on April 24, 1979. Horizontal bars indicate 1 $\sigma$  error bar. (From McCormick et al. 1982.)

inhomogeneous distribution of nitrogen dioxide resulting from its diurnal variation along the slant tangent path has been neglected in the retrieval of nitrogen dioxide vertical profiles. This effect has been




Figure 6. SAGE I measured ozone profile and coincident ECC ozonesonde profile obtained at Garmisch-Partenkirchen, Germany. Profiles obtained on April 2, 1979. ECC ozonesonde data starting at 1645 GMT; SAGE I measurements at 1743 GMT. (From Reiter and McCormick 1982.) Horizontal bars indicate 1 $\sigma$  error bar.




Figure 7. SAGE I sunset nitrogen dioxide data and previous measurements of nitrogen dioxide as shown in WMO (1981). Horizontal bars indicate 1 $\sigma$  error bar. (From Chu and McCormick 1986.)

analyzed by Kerr, Evans, and McConnell (1977) and the correction factors were found to be small. The SAGE II instrument uses two channels with relatively narrow bandwidths to make the nitrogen dioxide measurement, and this retrieval results in a significant improvement in the profile accuracy from that experiment (Cunnold et al. 1991).

SAGE I data were originally processed and archived at NSSDC in 1985, with corrections to NOAA temperature data at upper pressure levels as tabulated by Geller, Wu, and Gelman (1983). However, the tabulated correction factors were later determined to have sign errors (Gelman et al. 1986). The effects on the original SAGE I results were most pronounced in low-latitude-high-altitude regions. As part of the activities of the NASA-World Meteorological Organization (WMO) Ozone Trends Panel in 1987 (Watson et al. 1988), SAGE I data were reprocessed with the corrected temperature adjustment factors. This user's guide describes the revised SAGE I data set that has replaced the old SAGE I data set at NSSDC.

## Data Validation

Before being archived, the SAGE I data were validated through an extensive correlative measurements program. Two major correlative measurement experiments were conducted—the first over Natal, Brazil, in April 1979, and the second over Poker Flat Research Range, Alaska, in July 1979. In each experiment, correlative aerosol and ozone measurements were made nearly coincident in space and time with the SAGE I measurements. The correlative aerosol data consisted of SAM II-measured aerosol extinction profiles, airborne lidar-measured aerosol backscatter data, balloon-borne optical particle counter (dustsonde) data, and other in situ particle counter measurements. These correlative aerosol measurements were collected and converted to aerosol extinctions for comparison with SAGE I aerosol data (Russell, McCormick, et al. 1981; Russell, Swissler et al. 1981; Russell et al. 1984). The correlative ozone data consisted of ECC ozonesondes measurements and rocket ozone data (McCormick et al. 1984). Figure 8 shows the aerosol data from the Poker Flat experiment, along with the SAGE I, the SAM II, the dustsonde (DS), filter, and the Ames Wire Impactor (AWI) data (Russell et al. 1984). Other comparisons of the SAGE I and SAM II aerosol extinction measurements have also been made (Yue, McCormick, and Chu 1984). Figure 9 shows the SAGE I ozone measurements and the ECC ozonesonde and rocket measurements at Poker Flat and Wallops Island, Virginia (McCormick et al. 1989). Comparisons of the SAGE I ozone data with other ECC data have also been made (Reiter and McCormick 1982). Because of the unavailability of any generally accepted nitrogen dioxide monitor, the SAGE I nitrogen dioxide measurements were compared with available climatology as well as noncoincident balloon

soundings and ground-based measurements (Chu and McCormick 1986).

These comparisons between SAGE I aerosol, ozone, and nitrogen dioxide measurements and correlative data demonstrate that the SAGE I measurements agree with the correlative data to within their measurement and conversion uncertainties. These results clearly support the validity of the SAGE I aerosol, ozone, and nitrogen dioxide data.

## Data Products

The basic archived products from the SAGE I data processing are the aerosol profile tape, the ozone profile tape, and the nitrogen dioxide profile tape. These data products are archived and can be requested on magnetic tape from the National Space Sciences Data Center (NSSDC), NASA Goddard Space Flight Center, Code 933, Greenbelt, Maryland 20771. (Foreign requests should be sent to Code 930.2.) Each request should specify the experiment data desired (aerosol, ozone, or nitrogen dioxide), the NSSDC identification number (79-013A-01C), and the time period of interest. All 33 months of SAGE I data have been archived.

The aerosol profile tape contains the altitude profiles (plus the corresponding error estimates) of aerosol extinction coefficients at wavelengths of 1000 and 450 nm, Rayleigh extinction coefficients at wavelengths of 1000 and 450 nm, and the extinction ratio of aerosol extinction coefficient to Rayleigh extinction coefficient at a wavelength of 1000 nm. Auxiliary information that was used for processing the SAGE I data, such as NOAA's temperature-versus-altitude data, ephemeris-calculated latitude and longitude at the tangent location, beta angle of the spacecraft orbit (angle between the spacecraft orbit plane and the Earth-Sun direction), and event duration, is also included on the profile tape. There is one aerosol profile tape per year that covers 12 months of data from December of 1 year to November of the next year (winter to fall). The tapes are formatted as 6250-bpi, 9-track tapes and are archived at NSSDC. The aerosol profile tape record format is described in appendix A.

The ozone profile tapes contain the sunrise and sunset altitude profiles of ozone number density and mixing ratio, plus their corresponding error estimates. They contain the same auxiliary information as the aerosol profile tapes. There is one ozone profile tape per year that covers 12 months of data from December of 1 year to November of the next year. The tapes are formatted as 6250-bpi, 9-track tapes and are archived at NSSDC. The ozone profile tape format is described in appendix B.




Figure 8. SAGE I aerosol extinction coefficient retrieval and correlative aerosol measurements collected during 1979 validation experiment over Poker Flat Research Range, Alaska. Arrows in figure indicate height of tropopause.  $\lambda = 1.0 \mu\text{m}$ . DS = Dustsonde; AWI = Ames Wire Impactor. (From Russell et al. 1984.)




Figure 9. SAGE I ozone data with coincident ECC ozonesonde data and rocket ozonesonde data. Horizontal bars indicate  $1\sigma$  error bar. (From McCormick et al. 1989.)

The nitrogen dioxide profile tapes contain SAGE I nitrogen dioxide data in the same format as the ozone profile tapes. Similarly, there is one nitrogen dioxide data tape per year that covers 12 months of data from December of 1 year to November of the next year. The nitrogen dioxide profile tape format is described in appendix C.

Appendixes A, B, and C refer to the meteorological data notes of appendix D to explain the meteorological fields supplied by the National Meteorological Center in the data records of the tapes. Appendix D also explains how the meteorological data were handled by the SAGE I processing team.

## Scientific Studies

The SAGE I data have been used to perform a variety of atmospheric studies of atmospheric chemistry and dynamics, history of volcanic perturbation, and ozone and nitrogen dioxide long-term trends. The following discussion briefly describes some results of these investigations that used the SAGE I aerosol, ozone, and nitrogen dioxide data.

SAGE I aerosol data provided the first global description of volcanic perturbation of the stratospheric dust layer during the 33 months that SAGE I was in operation (Kent and McCormick 1984). Figure 10 shows the temporal variation of the optical depth of global stratospheric aerosols from 1979 to 1981, based on the combined SAGE I and SAM II aerosol measurements. It shows the effects of stratospheric aerosol loading resulting from the different volcanic

injections on both the Northern Hemisphere and the Southern Hemisphere. The two wavelengths of aerosol extinction coefficients obtained from the SAGE I measurements have been used to estimate the stratospheric aerosol size information, spatial distribution, and temporal evolution (Lenoble and Pruvost 1983; Yue and Deepak 1983; Yue and Deepak 1984).

The primary goal of the SAGE I mission was to monitor stratospheric trace species. However, the SAGE I instrument did make measurements in the troposphere at the cloud-free regions. Studies of the aerosols in the free troposphere, as measured by SAGE I, have been performed to research seasonal variability (Kent et al. 1988). Also, the SAGE I tropospheric data can be used to generate a high-altitude cloud climatology (Woodbury and McCormick 1986). Figure 11 illustrates a global map that was developed by the SAGE I data which shows the frequency of occurrence of the high-altitude clouds.

SAGE I aerosol, ozone, and nitrogen dioxide data have also been used for studies of atmospheric dynamics. For example, high-latitude aerosol, ozone, and nitrogen dioxide data have been used to study stratospheric sudden warming phenomena (Wang and McCormick 1985; Chu and McCormick 1986). SAGE I ozone data have been used to study the latitudinal transport in the atmosphere caused by wave activities (Wang, McCormick, and Chu 1983).




Figure 10. Temporal variation of optical depth of mean hemispheric stratospheric aerosols based on combined SAGE I and SAM II measurements from 1979 to 1981. Mass loading conversion factor =  $1.10 \times 10^3 \text{ m}^2 \text{ kg}^{-1}$ . (From Kent and McCormick 1984.)






Figure 11. Average frequency of cirrus cloud occurrence during full 33 months of SAGE I operation. (From Woodbury and McCormick 1986.)

SAGE I ozone and nitrogen dioxide data have also been used to study atmospheric chemistry anomalies related to nitrogen dioxide photochemistry (Chu and McCormick 1986) and ozone chemistry (Wang, McCormick, and Chu 1983).

The SAGE I ozone data have been compared with SAGE II ozone data to determine the global ozone trends from 1979 to late 1980 (Watson et al. 1988; McCormick, Veiga, and Zawodny 1989). It was this comparison between SAGE I and SAGE II ozone data that demonstrated stratospheric ozone at 50 km did not decrease significantly over the period studied contrary to the trend results based on other satellite observations (Herman, Hudson, and Serafino 1990; Chandra et al. 1990). The SAGE I and SAGE II ozone data have been regarded as the most accurate ozone profile data measured from space (WMO 1990).

## Concluding Remarks

This report has described the Stratospheric Aerosol and Gas Experiment I (SAGE I) instrument; its data collection, processing, and validation techniques; and the aerosol, ozone, and nitrogen dioxide archival products used in scientific investigations of various atmospheric sciences. These data products are archived and can be requested on magnetic tape from the National Space Sciences Data Center (NSSDC), NASA Goddard Space Flight Center, Code 933, Greenbelt, Maryland 20771. All 33 months of SAGE I data have been archived.

NASA Langley Research Center  
Hampton, VA 23681-0001  
May 28, 1992

## Appendix A

### SAGE I Aerosol Profile Record Format

#### Record Format

CYBER <sup>a</sup> words (60-bit)	Size	Field content description	Note (b)
40-km reference data			
0001	1	Event Date (yymmdd.0)	1
0002	↓	Event Time (hhmmss.0)	1
0003	↓	Subtangent Latitude ( $0.0^\circ \pm 90.0^\circ$ )	
0004	↓	Subtangent Longitude ( $0.0^\circ \pm 180.0^\circ$ )	
0005	↓	Spacecraft-Referenced Event Type (0.0 = Sunrise; 1.0 = Sunset)	2
0006	↓	Earth-Referenced Event Type (0.0 = Sunrise; 1.0 = Sunset)	2
0007	↓	Spacecraft Beta Angle ( $0.0^\circ \pm 61.0^\circ$ )	3
0008	↓	Coded Time of Year (ddd.fract)	4
NMC meteorological data (see appendix D)			
0009-0033	25	Temperature, K	
0034-0058	↓	Temperature Error, K	
0059-0083	↓	Geometric Altitude, m	
0084-0108	↓	Air Density, g/m <sup>3</sup>	
0109-0133	↓	Air Density Error, percent	
0134	1	Temperature Correction Value for 5.0-mbar Level, K	
0135	↓	Temperature Correction Value for 2.0-mbar Level, K	
0136	↓	Temperature Correction Value for 1.0-mbar Level, K	
0137	↓	Temperature Correction Value for 0.4-mbar Level, K	
0138	↓	"Meteorological Data Not Complete" Flag (0 = Complete; 1 = Incomplete)	
0139	↓	"Start of Model Meteorological Data" Array Index Pointer (1 19)	
0140	↓	Model Meteorological Data Selection Code (ssl)	
0141	↓	Revision Date of LaRC Meteorological Model (yymmdd.0)	
NASA LaRC processing information			
0142	1	LaRC Driver Revision Level	
0143	↓	LaRC Transmission Revision Level	
0144	↓	LaRC Inversion Revision Level	
0145	↓	LaRC Event Tag (yymmddhhmm.sq)	5
0146	↓	LaRC Processing Date (yymmdd.0)	1
0147	↓	LaRC Processing Time (hhmmss.0)	1
0148	↓	Mean Subtangent Altitude for Event Limb Calibration, km	6
0149	↓	<b>Value Designated as the Data Fill Number for this Event</b>	7
Event ground-track slew data			
0150-0157	8	Subtangent Altitude, km	
0158-0165	8	Corresponding Latitude ( $0.0^\circ \pm 90.0^\circ$ )	
0166-0173	8	Corresponding Longitude ( $0.0^\circ \pm 180.0^\circ$ )	
0174	1	Time Span of Data from Levels 1 to 70, sec	
Altitude and meteorological data for profile arrays			
0175-0244	70	Geometric Altitude, km	
0245-0314	70	Corresponding Pressure, mbar	
0315-0384	70	Corresponding Temperature, K	
0385-0390	6	Spare	

<sup>a</sup>Trademark of Control Data Corporation.

<sup>b</sup>Data field notes given after table.

CYBER <sup>a</sup> words (60-bit)	Size	Field content description	Note (b)	
Quality estimations of channel optical depth profile				
0391	1 ↓	1000-nm-Wavelength Quality Factor	8	
0392		Spare	8	
0393		Spare		
0394		Spare		
0395		450-nm-Wavelength Quality Factor		
0396		Spare		
0397		Spare		
0398-0400	3	Spare		
Rayleigh extinction profiles				
0401-0460	60 ↓	1000-nm Rayleigh Extinction, km <sup>-1</sup>		
0461-0520		1000-nm Rayleigh Extinction Error, km <sup>-1</sup>		
0521-0580		Spare		
0581-0640		Spare		
0641-0700		450-nm Rayleigh Extinction, km <sup>-1</sup>		
0701-0760		450-nm Rayleigh Extinction Error, km <sup>-1</sup>		
0761-0820		Spare		
0821-0880		Spare		
Aerosol profiles				
0881-0940	60 ↓	1000-nm Extinction, km <sup>-1</sup>		
0941-1000		1000-nm Extinction Error, km <sup>-1</sup>		
1001-1060		Spare		
1061-1120		Spare		
1121-1180		450-nm Extinction, km <sup>-1</sup>		
1181-1240		450-nm Extinction, km <sup>-1</sup>		
1241-1300		Spare		
1301-1360		Spare		
1361-1420		1000-nm Extinction Ratio		9
1421-1480		1000-nm Extinction Ratio Error		9
1481-1488	8	Spare		
End of event record				

<sup>a</sup>Trademark of Control Data Corporation.

<sup>b</sup>Data field notes given after table.

## Record Format Notes

### General notes:

- Each field of the event record contains one 60-bit CYBER<sup>a</sup> floating point number.
- All time and data references are to GMT, except fields 146 and 147, which are LaRC processing time.
- All latitudes and longitudes are given at the event subtangential point.
- If any field in the event record is considered invalid, or has missing data, a fill value will be placed in that field. For each event record, that fill value can be found in field 149. (See data field note 7 below.)
- Each profile level bin is centered at the 0.5-km point within a range of 1.0 km.

### Data field notes:

1. The "yymmdd.0" and "hhmmss.0" fields are generated by the respective FORTRAN statements:

DATE = FLOAT (IYY\*10000 + IMM\*100 + IDD)

TIME = FLOAT (IHH\*10000 + IMM\*100 + ISS)

2. Spacecraft-Referenced Event Type and Earth-Referenced Event Type fields are normally the same type, but if the absolute value of the spacecraft beta angle is close to 61°, their values may be opposite. The Earth-Referenced Event Type field is based on a ground-observer's viewpoint.
3. The Spacecraft Beta Angle field is defined as the angle generated by the intersection of the Earth-Sun vector and the spacecraft orbit plane.
4. The Coded Time of Year field is the time at the beginning of the event (not the same time as for fields 1 and 2) and is generated by the FORTRAN statement:

CODTIME = FLOAT(DOY) + (SOD/86400.0)

where

DOY = Day of year (1-366)

SOD = Seconds of the day (0.0-86399.99...)

5. The LaRC Event Tag field is generated with statements similar to those in note 1. The ".sq" at the end of the value is the event number of the day divided by 100.
6. The Mean Subtangential Altitude for Event Limb Calibration field contains the altitude at which data for the exoatmospheric solar image were gathered for use in solar limb normalization for the event.
7. The Value Designated as the Data Fill Number for This Event field must be used to determine what data in the event record are valid. If any field other than this one contains this number, that field has no valid information and should not be used by the investigator.
8. The Quality Factor fields for each wavelength are equal to 1.0 minus the summation of the optical depth errors at each profile level from 20.5 to 59.5 km. In cases where a 40-km span cannot be realized, the quality factor is proportioned to a 40-km span to allow a better comparison across wavelengths and other events.
9. Extinction ratio = (Aerosol extinction + Rayleigh extinction)/Rayleigh extinction.

<sup>a</sup>Trademark of Control Data Corporation.

## Appendix B

### SAGE I Ozone Profile Record Format

#### Record Format

CYBER <sup>a</sup> words (60-bit)	Size	Field content description	Note (b)
40-km reference data			
0001	1	Event Date (yymmdd.0)	1
0002	↓	Event Time (hhmmss.0)	1
0003	↓	Subtangent Latitude ( $0.0^\circ \pm 90.0^\circ$ )	
0004	↓	Subtangent Longitude ( $0.0^\circ \pm 180.0^\circ$ )	
0005	↓	Spacecraft-Referenced Event Type (0.0 = Sunrise; 1.0 = Sunset)	2
0006	↓	Earth-Referenced Event Type (0.0 = Sunrise; 1.0 = Sunset)	2
0007	↓	Spacecraft Beta Angle ( $0.0^\circ \pm 61.0^\circ$ )	3
0008	↓	Coded Time of Year (ddd.fract)	4
NMC meteorological data (see appendix D)			
0009-0033	25	Temperature, K	
0034-0058	↓	Temperature Error, K	
0059-0083	↓	Geometric Altitude, m	
0084-0108	↓	Air Density, $\text{g/m}^3$	
0109-0133	↓	Air Density Error, percent	
0134	1	Temperature Correction Value for 5.0-mbar Level, K	
0135	↓	Temperature Correction Value for 2.0-mbar Level, K	
0136	↓	Temperature Correction Value for 1.0-mbar Level, K	
0137	↓	Temperature Correction Value for 0.4-mbar Level, K	
0138	↓	"Meteorological Data Not Complete" Flag (0 = Complete; 1 = Incomplete)	
0139	↓	"Start of Model Meteorological Data" Array Index Pointer (1-19)	
0140	↓	Model Meteorological Data Selection Code (ssll)	
0141	↓	Revision Date of LaRC Meteorological Model (yymmdd.0)	
NASA LaRC processing information			
0142	1	LaRC Driver Revision Level	
0143	↓	LaRC Transmission Revision Level	
0144	↓	LaRC Inversion Revision Level	
0145	↓	LaRC Event Tag (yymmddhhmm.sq)	5
0146	↓	LaRC Processing Date (yymmdd.0)	1
0147	↓	LaRC Processing Time (hhmmss.0)	1
0148	↓	Mean Subtangent Altitude for Event Limb Calibration, km	6
0149	↓	<b>Value Designated as the Data Fill Number for this Event</b>	7
Event ground-track slew data			
0150-0157	8	Subtangent Altitude, km	
0158-0165	8	Corresponding Latitude ( $0.0^\circ \pm 90.0^\circ$ )	
0166-0173	8	Corresponding Longitude ( $0.0^\circ \pm 180.0^\circ$ )	
0174	1	Time Span of Data from Levels 1 to 70, sec	
Altitude and meteorological data for profile arrays			
0175-0244	70	Geometric Altitude, km	
0245-0314	70	Corresponding Pressure, mbar	
0315-0384	70	Corresponding Temperature, K	
0385-0390	6	Spare	

<sup>a</sup>Trademark of Control Data Corporation.

<sup>b</sup>Data field notes given after table.

CYBER <sup>a</sup> words (60-bit)	Size	Field content description	Note (b)
Quality estimations of channel optical depth profile			
0391	1 ↓	Spare	8
0392		Spare	
0393		600-nm-Wavelength Quality Factor	
0394		Spare	
0395		Spare	
0396		Spare	
0397		Spare	
0398-0400	3	Spare	
Ozone profiles			
0401-0470	70 ↓	Number Density, molecules/cm <sup>3</sup>	
0471-0540		Number Density Error, molecules/cm <sup>3</sup>	
0541-0610		Volumetric Mixing Ratio, v/v	
0611-0680		Volumetric Mixing Ratio Error, v/v	
0681-0688	8	Spare	
End of event record			

<sup>a</sup>Trademark of Control Data Corporation.

<sup>b</sup>Data field notes given after table.

## Record Format Notes

### General notes:

- Each field of the event record contains one 60-bit CYBER<sup>a</sup> floating point number.
- All time and data references are to GMT, except fields 146 and 147, which are LaRC processing time.
- All latitudes and longitudes are given at the event subangent point.
- If any field in the event record is considered invalid, or has missing data, a fill value will be placed in that field. For each event record, that fill value can be found in field 149. (See data field note 7 below.)
- Each profile level bin is centered at the 0.5-km point within a range of 1.0 km.

### Data field notes:

1. The "yymmdd.0" and "hhmmss.0" fields are generated by the respective FORTRAN statements:

DATE = FLOAT (IYY\*10000 + IMM\*100 + IDD)

TIME = FLOAT (IHH\*10000 + IMM\*100 + ISS)

2. Spacecraft-Referenced Event Type and Earth-Referenced Event Type fields are normally the same type, but if the absolute value of the spacecraft beta angle is close to 61°, their values may be opposite. The Earth-Referenced Event Type field is based on a ground-observer's viewpoint.
3. The Spacecraft Beta Angle field is defined as the angle generated by the intersection of the Earth-Sun vector and the spacecraft orbit plane.
4. The Coded Time of Year field is the time at the beginning of the event (not the same time as for fields 1 and 2) and is generated by the FORTRAN statement:

CODTIME = FLOAT(DOY) + (SOD/86400.0)

where

DOY = Day of year (1-366)

SOD = Seconds of the day (0.0-86399.99...)

5. The LaRC Event Tag field is generated with statements similar to those in note 1. The ".sq" at the end of the value is the event number of the day divided by 100.
6. The Mean Subangent Altitude for Event Limb Calibration field contains the altitude at which data for the exoatmospheric solar image were gathered for use in solar limb normalization for the event.
7. The Value Designated as the Data Fill Number for This Event field must be used to determine what data in the event record are valid. If any field other than this one contains this number, that field has no valid information and should not be used by the investigator.
8. The Quality Factor fields for each wavelength are equal to 1.0 minus the summation of the optical depth errors at each profile level from 20.5 to 59.5 km. In cases where a 40-km span cannot be realized, the quality factor is proportioned to a 40-km span to allow a better comparison across wavelengths and other events.

<sup>a</sup>Trademark of Control Data Corporation.

## Appendix C

### SAGE I Nitrogen Dioxide Profile Record Format

#### Record Format

CYBER <sup>a</sup> words (60-bit)	Size	Field content description	Note (b)
40-km reference data			
0001	1	Event Date (yymmdd.0)	1
0002	↓	Event Time (hhmmss.0)	1
0003	↓	Subtangent Latitude ( $0.0^\circ \pm 90.0^\circ$ )	
0004	↓	Subtangent Longitude ( $0.0^\circ \pm 180.0^\circ$ )	
0005	↓	Spacecraft-Referenced Event Type (0.0 = Sunrise; 1.0 = Sunset)	2
0006	↓	Earth-Referenced Event Type (0.0 = Sunrise; 1.0 = Sunset)	2
0007	↓	Spacecraft Beta Angle ( $0.0^\circ \pm 61.0^\circ$ )	3
0008	↓	Coded Time of Year (ddd.fract)	4
NMC meteorological data (see appendix D)			
0009-0033	25	Temperature, K	
0034-0058	↓	Temperature Error, K	
0059-0083	↓	Geometric Altitude, m	
0084-0108	↓	Air Density, $\text{g/m}^3$	
0109-0133	↓	Air Density Error, percent	
0134	1	Temperature Correction Value for 5.0-mbar Level, K	
0135	↓	Temperature Correction Value for 2.0-mbar Level, K	
0136	↓	Temperature Correction Value for 1.0-mbar Level, K	
0137	↓	Temperature Correction Value for 0.4-mbar Level, K	
0138	↓	"Meteorological Data Not Complete" Flag (0 = Complete; 1 = Incomplete)	
0139	↓	"Start of Model Meteorological Data" Array Index Pointer (1-19)	
0140	↓	Model Meteorological Data Selection Code (ssll)	
0141	↓	Revision Date of LaRC Meteorological Model (yymmdd.0)	
NASA LaRC processing information			
0142	1	LaRC Driver Revision Level	
0143	↓	LaRC Transmission Revision Level	
0144	↓	LaRC Inversion Revision Level	
0145	↓	LaRC Event Tag (yymmddhhmm.sq)	5
0146	↓	LaRC Processing Date (yymmdd.0)	1
0147	↓	LaRC Processing Time (hhmmss.0)	1
0148	↓	Mean Subtangent Altitude for Event Limb Calibration, km	6
0149	↓	<b>Value Designated as the Data Fill Number for this Event</b>	7
Event ground-track slew data			
0150-0157	8	Subtangent Altitude, km	
0158-0165	8	Corresponding latitude ( $0.0^\circ \pm 90.0^\circ$ )	
0166-0173	8	Corresponding Longitude ( $0.0^\circ \pm 180.0^\circ$ )	
0174	1	Time Span of Data from Levels 1 to 70, sec	
Altitude and meteorological data for profile arrays			
0175-0244	70	Geometric Altitude, km	
0245-0314	70	Corresponding Pressure, mbar	
0315-0384	70	Corresponding Temperature, K	
0385-0390	6	Spare	

<sup>a</sup>Trademark of Control Data Corporation.

<sup>b</sup>Data field notes given after table.



CYBER <sup>a</sup> words (60-bit)	Size	Field content description	Note (b)
Quality estimations of channel optical depth profile			
0391	1	Spare	8
0392	↓	Spare	
0393		Spare	
0394		Spare	
0395		450-nm-Wavelength Quality Factor	
0396		Spare	
0397	↓	385-nm-Wavelength Quality Factor	8
0398-0400	3	Spare	
Nitrogen dioxide profiles			
0401-0460	60	Number Density, molecules/cm <sup>3</sup>	
0461-0520	↓	Number Density Error, molecules/cm <sup>3</sup>	
0521-0580		Volumetric Mixing Ratio, v/v	
0581-0640		Volumetric Mixing Ratio Error, v/v	
End of event record			

<sup>a</sup>Trademark of Control Data Corporation.

<sup>b</sup>Data field notes given after table.

## Record Format Notes

### General notes:

- Each field of the event record contains one 60-bit CYBER<sup>a</sup> floating point number.
- All time and data references are to GMT, except fields 146 and 147, which are LaRC processing time.
- All latitudes and longitudes are given at the event sub-tangent point.
- If any field in the event record is considered invalid, or has missing data, a fill value will be placed in that field. For each event record, that fill value can be found in field 149. (See data field note 7 below.)
- Each profile level bin is centered at the 0.5-km point within a range of 1.0 km.

### Data field notes:

1. The "yymmdd.0" and "hhmmss.0" fields are generated by the respective FORTRAN statements:

$$\text{DATE} = \text{FLOAT} (\text{IYY} \times 10000 + \text{IMM} \times 100 + \text{IDD})$$
$$\text{TIME} = \text{FLOAT} (\text{IHH} \times 10000 + \text{IMM} \times 100 + \text{ISS})$$

2. Spacecraft-Referenced Event Type and Earth-Referenced Event Type fields are normally the same type, but if the absolute value of the spacecraft beta angle is close to 61°, their values may be opposite. The Earth-Referenced Event Type field is based on a ground-observer's viewpoint.
3. The Spacecraft Beta Angle field is defined as the angle generated by the intersection of the Earth-Sun vector and the spacecraft orbit plane.
4. The Coded Time of Year field is the time at the beginning of the event (not the same time as for fields 1 and 2) and is generated by the FORTRAN statement:

$$\text{CODTIME} = \text{FLOAT}(\text{DOY}) + (\text{SOD}/86400.0)$$

where

DOY = Day of year (1-366)

SOD = Seconds of the day (0.0-86399.99...)

5. The LaRC Event Tag field is generated with statements similar to those in note 1. The ".sq" at the end of the value is the event number of the day divided by 100.
6. The Mean Subtangent Altitude for Event Limb Calibration field contains the altitude at which data for the exoatmospheric solar image were gathered for use in solar limb normalization for the event.
7. The Value Designated as the Data Fill Number for This Event field must be used to determine what data in the event record are valid. If any field other than this one contains this number, that field has no valid information and should not be used by the investigator.
8. The Quality Factor fields for each wavelength are equal to 1.0 minus the summation of the optical depth errors at each profile level from 20.5 to 59.5 km. In cases where a 40-km span cannot be realized, the quality factor is proportioned to a 40-km span to allow a better comparison across wavelengths and other events.

<sup>a</sup>Trademark of Control Data Corporation.

## Appendix D

### Meteorological Data Notes

Meteorological data are supplied to NASA Langley Research Center (LaRC) by NOAA National Weather Service, Climate Analysis Branch, Washington, D.C. Data for temperature, temperature error, geometric altitude, air density, and air density error are provided for 18 pressure levels (1000 to 0.4 mbar) and for the derived tropopause pressure. The pressure levels (in mbar) correspond to the 25-element meteorological data arrays (1 to 25) as follows: 1000, 850, 700, 500, 400, 300, 250, 200, 150, 100, 70, 50, 30, 10, 5, 2, 1, 0.4, 0.04, 0.01, spare, spare, spare, spare, and derived tropopause pressure. Elements 19 and 20 (0.04 and 0.01 mbar) contain LaRC model data for temperature and altitude only. Elements 21 to 24 (spares) contain fill values, and element 25 contains the NOAA-supplied tropopause information.

If NOAA cannot supply meteorological data as above, LaRC determines the highest pressure level for which data are supplied and then inserts model data from the next level up to the lowest pressure level of 0.01 mbar. Only temperature and altitude information are supplied from using these model data. Temperature error, density, and density error will be filled for the corresponding levels that contain the LaRC-supplied data.

Meteorological correction factors for temperatures at 5, 2, 1, and 0.4 mbar are already added to the value of the temperatures in elements 15 through 18 of the temperature array. These correction factors are only included in the NOAA-supplied data. If model data are in these locations, no correction factors are applied. The correction factors are listed in fields 134 to 137 of the record.

Meteorological data (fields 245 to 384 of the event record) are interpolated from the meteorological data in fields 9 to 133 of the record. Altitude data in fields 175 to 244 increment by 1 km with the center of each level at the 0.5-km point of the level bin.

Other meteorological data information is contained in the following record locations:

Field 0138: 0 if NOAA-supplied data are complete, 1 if incomplete.

Field 0139: the model pointer is the array index that points to the start of LaRC-supplied model data in the temperature and altitude arrays of the meteorological data.

Field 0140: model selection code (ssll) where ss is from 01 to 04 for spring to winter, and ll is from 0 to 80 in 10-deg increments for absolute latitude.

Field 0141: date of revision of the LaRC-supplied model.

## References

- Chandra, S.; McPeters, R. D.; Hudson, R. D.; and Planet, W. 1990: Ozone Measurements From the NOAA-9 and the Nimbus-7 Satellites: Implications of Short and Long Term Variabilities. *Geophys. Res. Lett.*, vol. 17, no. 10, pp. 1573-1576.
- Chu, W. P.; and McCormick, M. P. 1979: Inversion of Stratospheric Aerosol and Gaseous Constituents From Spacecraft Solar Extinction Data in the 0.38-1.0- $\mu$ m Wavelength Region. *Appl. Opt.*, vol. 18, no. 9, pp. 1404-1413.
- Chu, W. P.; and McCormick, M. P. 1986: SAGE Observations of Stratospheric Nitrogen Dioxide. *J. Geophys. Res.*, vol. 91, no. D5, pp. 5465-5476.
- Chu, W. P.; McCormick, M. P.; Lenoble, J.; Brogniez, C.; and Pruvost, P. 1989: SAGE II Inversion Algorithm. *J. Geophys. Res.*, vol. 94, no. D6, pp. 8339-8351.
- Cunnold, D. M.; Zawodny, J. M.; Chu, W. P.; Pommereau, J. P.; and Goutail, F. 1991: Validation of SAGE II NO<sub>2</sub> Measurements. *J. Geophys. Res.*, vol. 96, pp. 12,913-12,925.
- Geller, Marvin A.; Wu, Mao-Fou; and Gelman, Melvyn E. 1983: Tropospheric-Stratosphere (Surface-55 km) Monthly Winter General Circulation Statistics for the Northern Hemisphere - Four Year Averages. *J. Atmos. Sci.*, vol. 40, no. 5, pp. 1334-1352.
- Gelman, M. E.; Miller, A. J.; Johnson, K. W.; and Nagatani, R. M. 1986: Detection of Long-Term Trends in Global Stratospheric Temperatures From NMC Analyses Derived From NOAA Satellite Data. *Adv. Space Res.*, vol. 6, no. 10, pp. 17-26.
- Herman, J. R.; Hudson, R. D.; and Serafino, G. 1990: Analysis of the Eight-Year Trend in Ozone Depletion From Empirical Models of Solar Backscattered Ultraviolet Instrument Degradation. *J. Geophys. Res.*, vol. 95, no. D6, pp. 7403-7416.
- Kent, G. S.; and McCormick, M. P. 1984: SAGE and SAM II Measurements of Global Stratospheric Aerosol Optical Depth and Mass Loading. *J. Geophys. Res.*, vol. 89, no. D4, pp. 5303-5314.
- Kent, G. S.; Farrukh, U. O.; Wang, P. H.; and Deepak, A. 1988: SAGE I and SAM II Measurements of 1  $\mu$ m Aerosol Extinction in the Free Troposphere. *J. Appl. Meteorol.*, vol. 27, pp. 269-279.
- Kerr, J. B.; Evans, W. F. J.; and McConnell, J. C. 1977: The Effects of NO<sub>2</sub> Changes at Twilight on Tangent Ray NO<sub>2</sub> Measurements. *Geophys. Res. Lett.*, vol. 4, no. 12, pp. 577-579.
- Lenoble, J.; and Pruvost, P. 1983: Inference of the Aerosol Angstrom Coefficient From SAGE Short-Wavelength Data. *J. Clim. & Appl. Meteorol.*, vol. 22, no. 10, pp. 1717-1725.
- McCormick, M. P.; Hamill, Patrick; Pepin, T. J.; Chu, W. P.; Swissler, T. J.; and McMaster, L. R. 1979: Satellite Studies of the Stratospheric Aerosol. *Bull. American Meteorol. Soc.*, vol. 60, no. 9, pp. 1038-1046.
- McCormick, M. P.; Kent, G. S.; Yue, G. K.; and Cunnold, D. M. 1982: Stratospheric Aerosol Effects From Soufrière Volcano as Measured by the SAGE Satellite System. *Science*, vol. 216, no. 4550, pp. 1115-1118.
- McCormick, M. P.; Swissler, T. J.; Hilsenrath, E.; Krueger, A. J.; and Osborn, M. T. 1984: Satellite and Correlative Measurements of Stratospheric Ozone: Comparison of Measurements Made by SAGE, ECC Balloons, Chemiluminescent, and Optical Rocketsondes. *J. Geophys. Res.*, vol. 89, no. D4, pp. 5315-5320.
- McCormick, M. P.; Zawodny, J. M.; Veiga, R. E.; Larsen, J. C.; and Wang, P. H. 1989: An overview of SAGE I and II Ozone Measurements. *Planet. & Space Sci.*, vol. 37, no. 12, pp. 1567-1586.
- McCormick, M. P.; Veiga, R. E.; and Zawodny, J. M. 1989: Comparison of SAGE I and SAGE II Stratospheric Ozone Measurements. *Ozone in the Atmosphere - Proceedings of the Quadrennial Ozone Symposium 1988 and Tropospheric Ozone Workshop*, Rumen D. Bojkov and Peter Fabian, eds., A. Deepak Publishing, pp. 202-205.
- McMaster, Leonard R. 1986: Stratospheric Aerosol and Gas Experiment (SAGE II). *Sixth Conference on Atmospheric Radiation*, American Meteorological Soc., pp. J46-J48.
- Reiter, R.; and McCormick, M. P. 1982: SAGE European Ozone-sonde Comparison. *Nature*, vol. 300, no. 5890, pp. 337-339.
- Russell, P. B.; McCormick, M. P.; Swissler, T. J.; Chu, W. P.; Livingston, J. M.; Fuller, W. H., Jr.; Rosen, J. M.; Hofmann, D. J.; McMaster, L. R.; Woods, D. C.; and Pepin, T. J. 1981: Satellite and Correlative Measurements of the Stratospheric Aerosol. II: Comparison of Measurements Made by SAM II, Dustsondes and an Airborne Lidar. *J. Atmos. Sci.*, vol. 38, no. 6, pp. 1295-1312.
- Russell, P. B.; Swissler, T. J.; McCormick, M. P.; Chu, W. P.; Livingston, J. M.; and Pepin, T. J. 1981: Satellite and Correlative Measurements of the Stratospheric Aerosol. I: An Optical Model for Data Conversions. *J. Atmos. Sci.*, vol. 38, no. 6, pp. 1279-1294.
- Russell, P. B.; McCormick, M. P.; Swissler, T. J.; Rosen, J. M.; Hofmann, D. J.; and McMaster, L. R. 1984: Satellite and Correlative Measurements of the Stratospheric Aerosol. III: Comparison of Measurements by SAM II, SAGE, Dustsondes, Filters, Impactors, and Lidar. *J. Atmos. Sci.*, vol. 41, no. 11, pp. 1791-1800.
- Wang, Pi-Huan; McCormick, M. P.; and Chu, W. P. 1983: A Study on the Planetary Wave Transport of Ozone During the Late February 1979 Stratospheric Warming Using the SAGE Ozone Observation and Meteorological Information. *J. Atmos. Sci.*, vol. 40, no. 10, pp. 2419-2431.
- Wang, Pi-Huan; and McCormick, M. P. 1985: Variations in Stratospheric Aerosol Optical Depth During Northern Warmings. *J. Geophys. Res.*, vol. 90, no. D6, pp. 10,597-10,606.

- Watson, R. T. et al. 1988: *Present State of Knowledge of the Upper Atmosphere 1988: An Assessment Report*. NASA RP-1208.
- WMO Global Ozone Research and Monitoring Project 1981: *The Stratosphere 1981—Theory and Measurements*. NASA TM-84125. (Available as WMO Rep. No. 11.)
- WMO Global Ozone Research and Monitoring Proj. 1990: *Scientific Assessment of Stratospheric Ozone: 1989, Volume I*. Rep. No. 20, World Meteorological Organization.
- Woodbury, Gerard E.; and McCormick, M. P. 1986: Zonal and Geographical Distributions of Cirrus Clouds Determined From SAGE Data. *J. Geophys. Res.*, vol. 91, no. D2, pp. 2775-2785.
- Yue, Glenn K.; and Deepak, Adarsh 1983: Retrieval of Stratospheric Aerosol Size Distribution From Atmospheric Extinction of Solar Radiation at Two Wavelengths. *Appl. Opt.*, vol. 22, no. 11, pp. 1639-1645.
- Yue, Glenn K.; and Deepak, Adarsh A. 1984: Latitudinal and Altitudinal Variation of Size Distribution of Stratospheric Aerosols Inferred From SAGE Aerosol Extinction Coefficient Measurements at Two Wavelengths. *Geophys. Res. Lett.*, vol. 11, no. 10, pp. 999-1002.
- Yue, Glenn K.; McCormick, M. P.; and Chu, W. P. 1984: A Comparative Study of Aerosol Extinction Measurements Made by the SAM II and SAGE Satellite Experiments. *J. Geophys. Res.*, vol. 89, no. D4, pp. 5321-5327.

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE August 1992	3. REPORT TYPE AND DATES COVERED Reference Publication		
4. TITLE AND SUBTITLE SAGE I Data User's Guide		5. FUNDING NUMBERS WU 665-45-30-21		
6. AUTHOR(S) L. R. McMaster, W. P. Chu, and M. W. Rowland				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) NASA Langley Research Center Hampton, VA 23681-0001		8. PERFORMING ORGANIZATION REPORT NUMBER L-16879		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) National Aeronautics and Space Administration Washington, DC 20546-0001		10. SPONSORING/MONITORING AGENCY REPORT NUMBER NASA RP-1275		
11. SUPPLEMENTARY NOTES McMaster and Chu: Langley Research Center, Hampton, VA; Rowland: ST Systems Corporation (STX), Hampton, VA.				
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Unclassified-Unlimited  Subject Category 46		12b. DISTRIBUTION CODE		
13. ABSTRACT (Maximum 200 words) This document is a guide for using the data products from the Stratospheric Aerosol and Gas Experiment I (SAGE I) for scientific investigations of stratospheric chemistry related to aerosol, ozone, nitrogen dioxide, dynamics, and climate change. Included is a detailed description of the aerosol profile tape, the ozone profile tape, and the nitrogen dioxide profile tape. These tapes are the SAGE I data products containing aerosol extinction data and ozone and nitrogen dioxide concentration data for use in the different scientific investigations. Also included are brief descriptions of the instrument operation; data collection, processing and validation; and some of the scientific analyses that have been conducted.				
14. SUBJECT TERMS SAGE; Aerosol; Ozone; Nitrogen dioxide		15. NUMBER OF PAGES 26		
		16. PRICE CODE A03		
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	